

Algorithmic randomness and monotone complexity on product space

Hayato Takahashi

The Institute of Statistical Mathematics
10-3 Midori-cho, Tachikawa, Tokyo 190-8562, Japan
hayato.takahashi@ieee.org

June 29, 2010

Abstract

We study algorithmic randomness and monotone complexity on product of the set of infinite binary sequences. We explore the following problems: monotone complexity on product space, Lambalgen's theorem for correlated probability, classification of random sets by likelihood ratio tests, decomposition of complexity and independence, Bayesian statistics for individual random sequences. Formerly Lambalgen's theorem for correlated probability is shown under a uniform computability assumption in [H. Takahashi Inform. Comp. 2008]. In this paper we show the theorem without the assumption.

Keywords : Martin-Löf randomness, Kolmogorov complexity, Lambalgen's Theorem, consistency, Bayesian statistics

1 Introduction

It is known that Martin-Löf random sequences [11] satisfy many laws of probability one, for example ergodic theorem, martingale convergence theorem, and so on, see [24, 19]. In this paper, we study Martin-Löf random sequences with respect to a probability on product space $\Omega \times \Omega$, where Ω is the set of infinite binary sequences. In particular, we investigate the following problems:

1. Randomness and monotone complexity on product space (Levin-Schnorr theorem for product space)
2. Lambalgen's theorem [22] for correlated probability.
3. Likelihood ratio test and classification of random sets.
4. Decomposition of complexity and independence of individual random sequences.
5. Bayesian statistics for individual random sequences.

The above problems are property of product space except for 3.

In Section 3, we show Lambalgen's theorem for correlated probability. In the previous paper [19], the theorem is shown under a uniform computability assumption. In this paper, we show the theorem without that assumption. This is the main theorem of this paper (Theorem 3.3).

The other sections are as follows: In Section 2, we define monotone complexity on product space. A usual definition of one-dimensional monotone complexity strongly depends on an order structure of one-dimensional space. In order to define monotone complexity on product space, we give an algebraic definition of monotone function for product space, which is applicable, *mutatis mutandis*, to an abstract partially ordered set. In Section 4, we show a classification of random sets by likelihood ratio tests. In particular we show an important theorem by Martin-Löf, i.e., two computable probabilities are mutually singular iff their random sets are disjoint. As a simple application, we show consistency of MDL for individual sequences. In Section 5, we show a decomposition of monotone complexity for prefixes of random sequences under a condition. As a corollary, we show some equivalent conditions for independence of individual random sequences. In Section 6, we apply our results to Bayesian statistics. By virtue of randomness theory, we can develop a point-wise theory for Bayesian statistics. In particular, we show consistency of posterior distribution (and its equivalent conditions) for individual random sequences. In order to show this, the results of Section 4 plays an important role. Also we show an asymptotic theory of estimation for individual sequences, which is closely related to decomposition of complexity.

2 Randomness and complexity

First we introduce Martin-Löf randomness on Ω . Let S be the set of finite binary strings. Let Ω be the set of infinite binary sequences with product topology. As in [19], we write $A \subset B$ including $A = B$. Throughout the paper, the base of logarithm is 2. We use symbols such as x, y, s to denote an element of S and x^∞, y^∞ to denote an element of Ω . For $x \in S$, let $\Delta(x) := \{x\omega : \omega \in \Omega\}$, where $x\omega$ is the concatenation of x and ω , and for $x^\infty \in \Omega$, $\Delta(x^\infty) := \{x^\infty\}$. Let $\lambda \in S$ be the empty word, then $\Delta(\lambda) = \Omega$. For $A \subset S$, let $\sigma\{\Delta(x)\}_{x \in A}$ be the σ -algebra generated by $\{\Delta(x)\}_{x \in A}$ and $\mathcal{B} := \sigma\{\Delta(x)\}_{x \in S}$. Let $(\Omega, \mathcal{B}, P)$ be a probability space. We write $P(x) := P(\Delta(x))$ for $x \in S$, then we have $P(x) = P(x0) + P(x1)$ for all x . Let $\mathbb{N}, \mathbb{Q}$, and $\mathbb{R}$ be the set of natural numbers, rational numbers, and real numbers, respectively. P is called computable if there exists a computable function $p : S \times \mathbb{N} \rightarrow \mathbb{Q}$ such that $\forall x \in S \forall k \in \mathbb{N} |P(x) - p(x, k)| < 1/k$. A set $A \subset S$ is called recursively enumerable (r.e.) if there is a computable function $f : \mathbb{N} \rightarrow S$ such that $f(\mathbb{N}) = A$. For $A \subset S$, let $\tilde{A} := \bigcup_{x \in A} \Delta(x)$. A set $U \subset \mathbb{N} \times S$ is called (Martin-Löf) test with respect to P if 1) U is r.e., 2) $\tilde{U}_{n+1} \subset \tilde{U}_n$ for all n , where $U_n = \{(n, x) \in U\}$, and 3) $P(\tilde{U}_n) < 2^{-n}$. In the following, if P is obvious from the context, we say that U is a test. A test U is called universal if for any other test V , there is a constant c such that $\forall n \tilde{V}_{n+c} \subset \tilde{U}_n$.

Theorem 2.1 (Martin-Löf[11]) *If P is a computable probability, a universal test U exists.*

In [11], the set $(\bigcap_{n=1}^{\infty} \tilde{U}_n)^c$ (complement of the limit of universal test) is defined to be random sequences with respect to P , where U is a universal test. We write $\mathcal{R}^P := (\bigcap_{n=1}^{\infty} \tilde{U}_n)^c$. Note that for two universal tests U and V , $\bigcap_{n=1}^{\infty} \tilde{U}_n = \bigcap_{n=1}^{\infty} \tilde{V}_n$ and hence $\mathcal{R}^P$ does not depend on the choice of a universal test. An equivalent definition of test is that U is r.e. and $\sum_n P(\tilde{U}_n) < \infty$. Then the set covered by $\tilde{U}_n$ infinitely many times is a limit of a test, i.e., $\limsup_n \tilde{U}_n \subset (\mathcal{R}^P)^c$, see [17].

For $x, y \in S$, let $\Delta(x, y) := \Delta(x) \times \Delta(y)$. Let $\mathcal{B}_{S^2} := \sigma\{\Delta(x, y) | x, y \in S\}$. Then computability of P on $(\Omega^2, \mathcal{B}_{S^2})$, its Martin-Löf tests, and the set of random sequences are defined similarly.

2.1 Complexity

For $x', x \in S \cup \Omega$, we write $x' \sqsubseteq x \Leftrightarrow x'$ is a prefix of $x \Leftrightarrow \Delta(x') \supset \Delta(x)$, and for $(x', y'), (x, y) \in (S \cup \Omega)^2$, $(x', y') \sqsubseteq (x, y) \Leftrightarrow x' \sqsubseteq x$ and $y' \sqsubseteq y \Leftrightarrow \Delta(x', y') \supset \Delta(x, y)$. Then $S \cup \Omega$ and $(S \cup \Omega)^2$ are partially ordered sets. For $A \subset S^2$, let $\bigvee A$ be the least upper bound of A . Then $\bigvee A$ exists in $(S \cup \Omega)^2$ iff $\bigcap_{(x,y) \in A} \Delta(x, y) \neq \emptyset$. In the following bold-faced symbols $\mathbf{x}, \mathbf{y}, \mathbf{p}$ denote an element of $(S \cup \Omega)^2$, $\mathbf{x}^\infty$ denote an element of Ω^2 , and $\boldsymbol{\lambda} = (\lambda, \lambda)$.

First we define monotone functions $(S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$. Let $F \subset S^2 \times S^2$ and $F_{\mathbf{p}} := \{\mathbf{x} | (\mathbf{p}, \mathbf{x}) \in F\}$.

Assume that

$$\forall \mathbf{p} \in S^2, \boldsymbol{\lambda} \in F_{\mathbf{p}} \text{ and } \bigvee_{\mathbf{p}' \sqsubseteq \mathbf{p}} F_{\mathbf{p}'} \text{ exists.} \quad (1)$$

Set

$$f(\mathbf{p}) := \bigvee_{\mathbf{p}' \sqsubseteq \mathbf{p}, \mathbf{p}' \in S^2} F_{\mathbf{p}'} \text{ for } \mathbf{p} \in (S \cup \Omega)^2. \quad (2)$$

We see that $f : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$ and f is monotone, i.e.,

$$\mathbf{p}' \sqsubseteq \mathbf{p} \Rightarrow f(\mathbf{p}') \sqsubseteq f(\mathbf{p}).$$

Conversely, let $f : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$ be a monotone function, and set

$$F := \{(\mathbf{p}, \mathbf{x}) \in S^2 \times S^2 | \mathbf{x} \sqsubseteq f(\mathbf{p})\},$$

Then $\bigvee F_{\mathbf{p}} = f(\mathbf{p})$, F satisfies (1), and the function defined by F coincides with f . If F is a r.e. set, the function f defined by (2) is called *computable monotone function*.

For $s \in S$, let $|s|$ be the length of s . In particular $|\lambda| = 0$ and $|x^\infty| = \infty$. For $\mathbf{p} = (p_1, p_2) \in (S \cup \Omega)^2$, let $|\mathbf{p}| := |p_1| + |p_2|$. The monotone complexity with respect to a computable monotone function $f : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$ is defined as follows:

$$Km_f^2(x, y) := \min\{|p_1| + |p_2| \mid (x, y) \sqsubseteq f(p_1, p_2)\},$$

$$Km_f(x, y) := \min\{|p| \mid (x, y) \sqsubseteq f(p, \lambda)\},$$

for $x, y, p, p_1, p_2 \in S \cup \Omega$. If there is no (p_1, p_2) such that $(x, y) \sqsubseteq f(p_1, p_2)$, then $Km_f^2(x, y) := \infty$. Similarly, $Km_f(x, y) := \infty$ if there is no p such that $(x, y) \sqsubseteq f(p, \lambda)$.

A computable monotone function $u : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$ is called *optimal* if for any computable monotone function $f : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$, there is a constant c such that $Km_u^2(\mathbf{x}) \leq Km_f^2(\mathbf{x}) + c$ for all $\mathbf{x} \in (S \cup \Omega)^2$. We can construct an optimal function in the following manner. First, observe that there is a r.e. set $\bar{F} \subset \mathbb{N} \times S^2 \times S^2$ such that 1) $F_i = \{(\mathbf{p}, \mathbf{x}) | (i, \mathbf{p}, \mathbf{x}) \in \bar{F}\}$ satisfies (1) for all $i \in \mathbb{N}$, and 2) for each r.e. set F that satisfies (1), there is i such that $F = F_i$. Note that the first condition in (1) is necessary to enumerate $\{F_i\}$. Next, set $F^u := \{(\bar{i}\mathbf{p}, \mathbf{x}) | (i, \mathbf{p}, \mathbf{x}) \in \bar{F}\}$, where $\bar{i}\mathbf{p} = (0^i 1 p_1, p_2)$ for $\mathbf{p} = (p_1, p_2)$. Let u be a computable monotone function defined by F^u via (2), then we see that u is optimal. In the following discussion, we fix u and let

$$\begin{aligned} Km^2(x, y) &:= Km_u^2(x, y), \quad Km(x, y) := Km_u(x, y), \\ Km(x|y) &:= \min\{|p| \mid (x, \lambda) \sqsubseteq u(p, y)\}, \\ Km(x) &:= Km(x|\lambda) \text{ for } x, y \in S \cup \Omega. \end{aligned}$$

By definition, we have $\forall x, y, \quad Km^2(x, y) \leq Km(x, y)$. Note that Km is equivalent to a monotone complexity that is defined from an optimal monotone function $S \cup \Omega \rightarrow (S \cup \Omega)^2$. Also note that $Km(x)$ defined above is different from $Km^2(x) := Km^2(x, \lambda)$. Later we show that Km^2 and Km are asymptotically bounded for prefixes of random sequences under a condition, see Corollary 2.1.

In the following, a subset $\mathcal{A}$ of $S \cup \Omega$ or $(S \cup \Omega)^2$ is called *non-overlapping* if $\Delta(\mathbf{x}) \cap \Delta(\mathbf{y}) = \emptyset$ for $\mathbf{x}, \mathbf{y} \in \mathcal{A}, \mathbf{x} \neq \mathbf{y}$. Note that $\Delta(\mathbf{x}) \cap \Delta(\mathbf{y}) = \emptyset \Rightarrow \mathbf{x}$ and $\mathbf{y}$ are incomparable. The converse is true if $\mathbf{x}, \mathbf{y} \in S \cup \Omega$. However if $\mathbf{x}, \mathbf{y} \in (S \cup \Omega)^2$ then there is a counter-example, e.g., $(\lambda, 0)$ and $(0, \lambda)$ are incomparable but $\Delta(\lambda, 0) \cap \Delta(0, \lambda) = \Delta(0, 0)$. In one-dimensional case, the notion of non-overlapping is equivalent to that of prefix-free. Throughout the paper we use the term “non-overlapping”.

Proposition 2.1 *a) monotonicity: $x \sqsubseteq z \Rightarrow Km(x|y) \leq Km(z|y)$, and $y \sqsubseteq z \Rightarrow Km(x|y) \geq Km(x|z)$.
b) Kraft inequality: $\sum_{\mathbf{x} \in \mathcal{A}} 2^{-Km(\mathbf{x})} \leq \sum_{\mathbf{x} \in \mathcal{A}} 2^{-Km^2(\mathbf{x})} \leq 1$ for non-overlapping set $\mathcal{A} \subset (S \cup \Omega)^2$.
c) Conditional sub-additivity: $\exists c \forall x, y \in S \cup \Omega, \quad Km^2(x, y) \leq Km(x|y) + Km(y) + c$.*

Proof) a) Obvious. b) Let u be an optimal monotone function and $p_{\mathbf{x}} \in \{\mathbf{p} | \mathbf{x} \sqsubseteq u(\mathbf{p})\}$. Suppose that $\Delta(\mathbf{x}) \cap \Delta(\mathbf{x}') = \emptyset$ and $\exists \mathbf{z}, \mathbf{z} = p_{\mathbf{x}} \vee p_{\mathbf{x}'}$. Then $\mathbf{x} \sqsubseteq$

$u(\mathbf{z})$ and $\mathbf{x}' \sqsubseteq u(\mathbf{z})$, which contradicts to $\Delta(\mathbf{x}) \cap \Delta(\mathbf{x}') = \emptyset$. Thus $\{p_{\mathbf{x}} | \mathbf{x} \in \mathcal{A}\}$ is non-overlapping for a non-overlapping set $\mathcal{A}$. By setting $p_{\mathbf{x}}$ to be an optimal code, i.e., $|p_{\mathbf{x}}| = Km_u^2(\mathbf{x})$, we have $\sum_{\mathbf{x} \in \mathcal{A}} 2^{-Km^2(\mathbf{x})} \leq 1$. Since $Km^2 \leq Km$, we have the statement. c) Let u be an optimal monotone function. Suppose that $x \sqsubseteq u(p, y)$, $Km(x|y) = |p|$ and $y \sqsubseteq u(p')$, $Km(y) = |p'|$. Let $f : (S \cup \Omega)^2 \rightarrow (S \cup \Omega)^2$ such that $f(p_1, p_2) := (u(p_1, u(p_2)), u(p_2))$ for all p_1, p_2 . Then f is monotone and $Km_f^2(x, y) \leq |p| + |p'| = Km(x|y) + Km(y)$. ■

Next we show Levin-Schnorr theorem for product space. Let $\mathcal{A} \subset S^2$ be a r.e. set and

$$\mathcal{A}(\mathbf{x}^\infty) := \{\mathbf{x} \in \mathcal{A} \mid \mathbf{x} \sqsubset \mathbf{x}^\infty\} \text{ for } \mathbf{x}^\infty \in \Omega^2.$$

Before proving the theorem, we need conditions on $\mathcal{A}$:

$$\mathbf{x}, \mathbf{y} \in \mathcal{A} \Rightarrow \mathbf{x} \text{ and } \mathbf{y} \text{ are comparable or } \Delta(\mathbf{x}) \cap \Delta(\mathbf{y}) = \emptyset. \quad (3)$$

If (3) holds then for any $\mathcal{A}' \subset \mathcal{A}$ there is a non-overlapping $\mathcal{A}'' \subset \mathcal{A}'$ such that $\tilde{\mathcal{A}}'' = \tilde{\mathcal{A}}'$. Note that it is possible $\mathcal{A}''$ is not r.e. even if $\mathcal{A}'$ is a r.e. set.

$$\mathbf{x}, \mathbf{y} \in \mathcal{A} \Rightarrow \exists \text{ non-overlapping } \alpha \subset \mathcal{A}, \Delta(\mathbf{x}) \cap (\Delta(\mathbf{y}))^c = \tilde{\alpha}. \quad (4)$$

Lemma 2.1 *If $\mathcal{A}$ is r.e. and satisfies (4) then for any r.e. $\mathcal{A}' \subset \mathcal{A}$ there is a non-overlapping r.e. $\mathcal{A}'' \subset \mathcal{A}$ such that $\tilde{\mathcal{A}}' = \tilde{\mathcal{A}}''$.*

Proof) Since $\mathcal{A}'$ is r.e., there is a computable $a' : \mathbb{N} \rightarrow \mathcal{A}'$ such that $a'(\mathbb{N}) = \mathcal{A}'$. Let $\mathcal{A}''(0) = \emptyset$. Suppose that $\mathcal{A}''(n-1)$ is a finite non-overlapping subset of $\mathcal{A}$ and $\tilde{\mathcal{A}}''(n-1) = \cup_{1 \leq i \leq n-1} \Delta(a'(i))$. Since $\mathcal{A}''(n-1)$ is finite, from (4), there is a non-overlapping $\alpha(n)$ such that

$$\tilde{\alpha}(n) = \Delta(a'(n)) \cap (\tilde{\mathcal{A}}''(n-1))^c. \quad (5)$$

Since $\Delta(a'(n)) \cap (\tilde{\mathcal{A}}''(n-1))^c$ is compact and $\alpha(n)$ is non-overlapping, from Heine-Borel Theorem, we see that $\alpha(n)$ is finite. Let $\beta(n) := \{\mathbf{z} \in \mathcal{A} \mid \Delta(\mathbf{z}) \subset \Delta(a'(n)) \cap (\tilde{\mathcal{A}}''(n-1))^c\}$. Since $\mathcal{A}$ is r.e. and $\mathcal{A}''(n-1)$ is finite, $\beta(n)$ is r.e. from $a'(n)$ and $\mathcal{A}''(n-1)$. In particular, since $\alpha(n) \subset \beta(n)$, we can compute a finite non-overlapping $\alpha(n)$ that satisfies (5) from $a'(n)$ and $\mathcal{A}''(n-1)$. Let $\mathcal{A}''(n) := \mathcal{A}''(n-1) \cup \alpha(n)$ then $\mathcal{A}''(n)$ is a finite non-overlapping set. Let $\mathcal{A}'' := \cup_n \mathcal{A}''(n)$. By induction, $\mathcal{A}'' \subset \mathcal{A}$ is a non-overlapping r.e. set such that $\tilde{\mathcal{A}}' = \tilde{\mathcal{A}}''$. ■

Theorem 2.2 (Levin-Schnorr theorem [9, 15, 16] on product space)

Let P be a computable probability on $(\Omega^2, \mathcal{B}_{S^2})$. Let $\mathcal{A}$ be a r.e. set that satisfies (3) and (4). Then

$$\mathbf{x}^\infty \in \mathcal{R}^P \Leftarrow \sup_{\mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty)} -\log P(\mathbf{x}) - Km(\mathbf{x}) < \infty, \mathbf{x}^\infty = \vee \mathcal{A}(\mathbf{x}^\infty).$$

$$\mathbf{x}^\infty \in \mathcal{R}^P \Rightarrow \sup_{\mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty)} -\log P(\mathbf{x}) - Km(\mathbf{x}) < \infty.$$

The above statements hold for Km^2 .

Proof) Suppose that $\mathbf{x}^\infty \notin \mathcal{R}^P$ and $\mathbf{x}^\infty = \vee \mathcal{A}(\mathbf{x}^\infty)$. Then there is a test U such that for all n , $\mathbf{x}^\infty \in \tilde{U}_n$ and $P(\tilde{U}_n) < 2^{-n}$. Let $U'_n := \{\mathbf{y} \in \mathcal{A} \mid \exists \mathbf{x} \in U_n, \mathbf{x} \subseteq \mathbf{y}\}$. Since U_n and $\mathcal{A}$ are r.e. sets, $U'_n \subset \mathcal{A}$ is a r.e. set. From Lemma 2.1, there is a non-overlapping r.e. set $U''_n \subset \mathcal{A}$ such that $\tilde{U}'_n = \tilde{U}''_n$. Since $\mathbf{x}^\infty = \vee \mathcal{A}(\mathbf{x}^\infty)$, we have $\mathbf{x}^\infty \in \tilde{U}''_n$ and $\forall n, U''_n \cap \mathcal{A}(\mathbf{x}^\infty) \neq \emptyset$. Let P' be a measure such that $P'(\mathbf{x}) = P(\mathbf{x})2^n$ for $\mathbf{x} \in U''_n$ and 0 otherwise. Since $P(\tilde{U}''_n) < 2^{-n}$, we have $\sum_{\mathbf{x} \in U''_n} P'(\mathbf{x}) < 1$. By applying Shannon-Fano-Elias coding to P' on U''_n , we have $\exists c_1, c_2 > 0 \forall n \exists \mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty) Km(\mathbf{x}) \leq -\log P(\mathbf{x}) - n + K(n) + c_1 \leq -\log P(\mathbf{x}) - n + 2 \log n + c_2$, where K is the prefix complexity.

Conversely, let $U_n := \{\mathbf{x} \in \mathcal{A} \mid Km(\mathbf{x}) < -\log P(\mathbf{x}) - n\}$. From (3), we see that there is a non-overlapping set $U'_n \subset U_n$ such that $\tilde{U}'_n = \tilde{U}_n$. Hence $P(\tilde{U}_n) = P(\tilde{U}'_n) < \sum_{\mathbf{x} \in U'_n} 2^{-Km(\mathbf{x})-n} \leq 2^{-n}$, where the last inequality follows from Proposition 2.1 b. Since U_n is a r.e. set, $\{U_n\}$ is a test and $\cap_n \tilde{U}_n \subset (\mathcal{R}^P)^c$. The proof for Km^2 is the same as above. ■

Example 1 Let $g : \mathbb{N} \rightarrow \mathbb{N}$ be a total-computable monotonically increasing function, where $n \leq m \Rightarrow g(n) \leq g(m)$. Let

$$\mathcal{A}_g := \{(x, y) \in S^2 \mid |y| = g(|x|)\}. \quad (6)$$

Then $\mathcal{A}_g$ is decidable and satisfies (3) and (4). If g is unbounded then $\forall \mathbf{x}^\infty, \vee \mathcal{A}_g(\mathbf{x}^\infty) = \mathbf{x}^\infty$.

Next we study a coding problem for multi-dimensional monotone complexity. The following lemma shows that if $\mathcal{A}$ is decidable and satisfies (3), we have the same one-dimensional coding as in [20].

Lemma 2.2 *Let P be a computable probability on $(\Omega^2, \mathcal{B}_{S^2})$ and let $\mathcal{A} \subset S^2$ be a decidable set that satisfies (3), then there is a computable monotone function $g : S \cup \Omega \rightarrow (S \cup \Omega)^2$ such that*

$$\exists c \forall \mathbf{x} \in A(\mathbf{x}^\infty), Km_g(\mathbf{x}) \leq -\log P(\mathbf{x}) + c.$$

Proof) If $\mathcal{A}$ is decidable and satisfies (3) then, by rearranging an enumeration of $\mathcal{A}$, we see that there is a computable $f : \mathbb{N} \rightarrow S^2$ such that $f(\mathbb{N}) = \mathcal{A}$ and $\forall i, j, i < j, \Delta(f(i)) \cap \Delta(f(j)) = \emptyset$ or $f(i) \sqsubseteq f(j)$. Then we can construct a family of half-open intervals $V_{f(i)} := [a(i), b(i)) \subset [0, 1], \forall i \in \mathbb{N}$ that satisfies the following conditions: 0) $V_\lambda = [0, 1], 1) |V_{f(i)}| = P(f(i))$ for all i , where $|V|$ is the length of the interval V , 2) if $\Delta(f(i)) \cap \Delta(f(j)) = \emptyset$ then $V_{f(i)} \cap V_{f(j)} = \emptyset$, 3) if $f(i) \sqsubseteq f(j)$ then $V_{f(i)} \supset V_{f(j)}$, and 4) a and b are computable, i.e., there are rational valued computable functions $A : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{Q}$ and $B : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{Q}$ such that $\forall i, k, |a(i) - A(i, k)| < 1/k, |b(i) - B(i, k)| < 1/k$. For $s = s_1 s_2 \cdots s_n \in S, \forall i, s_i \in \{0, 1\}$, let $I_s := [\sum_{1 \leq i \leq n} s_i 2^{-i}, \sum_{1 \leq i \leq n} s_i 2^{-i} + 2^{-n})$. Then set $F := \{(s, f(i)) \in S \times S^2 \mid I_s \subset V_{f(i)}, i \in \mathbb{N}\}$. We see that F is a r.e. set that satisfies (1). Let g be a computable monotone function defined by F , then we have $g : S \cup \Omega \rightarrow (S \cup \Omega)^2$ and $\exists c \forall \mathbf{x} \in A(\mathbf{x}^\infty), Km_g(\mathbf{x}) \leq -\log P(\mathbf{x}) + c$. $\blacksquare$

From Theorem 2.2, Lemma 2.2, and Proposition 2.1 c, we have

Corollary 2.1 *Let P be a computable probability on $(\Omega^2, \mathcal{B}_{S^2})$. If $\mathcal{A} \subset S^2$ is decidable and satisfies (3) and (4), then*

$$\begin{aligned} \mathbf{x}^\infty \in \mathcal{R}^P &\Rightarrow \sup_{\mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty)} |\log P(\mathbf{x}) + Km(\mathbf{x})| < \infty, \\ \mathbf{x}^\infty \in \mathcal{R}^P &\Leftarrow \sup_{\mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty)} |\log P(\mathbf{x}) + Km(\mathbf{x})| < \infty, \mathbf{x}^\infty = \vee \mathcal{A}(\mathbf{x}^\infty). \end{aligned}$$

The above statements are true for Km^2 , and

$$\begin{aligned} \mathbf{x}^\infty \in \mathcal{R}^P &\Rightarrow \sup_{\mathbf{x} \in \mathcal{A}(\mathbf{x}^\infty)} |Km(\mathbf{x}) - Km^2(\mathbf{x})| < \infty \\ &\Rightarrow \sup_{(x, y) \in \mathcal{A}(\mathbf{x}^\infty)} Km(x, y) - Km(x|y) - Km(y) < \infty. \end{aligned}$$

For 1-dimensional monotone complexity and its relation to other complexities, see [10, 21]. In [5], a conditional complexity K_* that is monotone with the conditional argument is defined.

Remark 1 It is not difficult to develop monotone function and complexity in an abstract way. Indeed, let A and $\bar{A}$ be partially ordered sets such that A is r.e. and $\bar{A} := \{\vee B | B \subset A\}$. Let $F \subset A \times A$ be a r.e. set that satisfies (1) with respect to the partial order of A . Then we can define (optimal) monotone function $f : \bar{A} \rightarrow \bar{A}$ in a similar way with Section 2.1. For example, for $\mathbf{x}, \mathbf{y} \in (S \cup \Omega)^\infty$, let $\mathbf{x} \sqsubseteq \mathbf{y}$ if $\forall i, x^i \sqsubseteq y^i$ for $\mathbf{x} = (x^1, x^2, \dots), \mathbf{y} = (y^1, y^2, \dots), x^i, y^i \in S \cup \Omega$. Then $(S \cup \Omega)^\infty$ is a partially ordered set. Let $A := \{(\mathbf{x}, \lambda^\infty) | \mathbf{x} \in \cup_k S^k\}$, where $\lambda^\infty = (\lambda, \lambda, \dots) \in S^\infty$. Then A is a sub-partially ordered set of $(S \cup \Omega)^\infty$ and $\bar{A} = (S \cup \Omega)^\infty$. We can define computable monotone function $f : \bar{A} \rightarrow \bar{A}$. For $\mathbf{x} = (x_1, \dots, x_n, \dots) \in \bar{A}$, let $|\mathbf{x}| := \sum_n |x_n|$. Then Km_f is defined. For example, let us consider discrete time (computable) stochastic processes $X_i \in \Omega, i = 1, 2, \dots$. Then their randomness and complexity of sample paths are modeled with a computable probability on $(\Omega^\infty, \mathcal{B}_A)$ and Km_f , where $\mathcal{B}_A := \sigma\{\Delta(\mathbf{x}) | \mathbf{x} \in A\}, \Delta(\mathbf{x}) := \{\mathbf{x}^\infty | \mathbf{x} \sqsubset \mathbf{x}^\infty \in \Omega^\infty\}$ and computability of probabilities on $(\Omega^\infty, \mathcal{B}_A)$ is defined in a similar manner with finite dimensional case.

Remark 2 Let $\phi^{l,t} : (S \cup \Omega)^l \rightarrow (S \cup \Omega)^t$ be an optimal monotone function for $1 \leq l, t \leq \infty$. Then $Km^{l,t}(x_1, \dots, x_t) \leq Km^{l',t}(x_1, \dots, x_t) + O(1)$ if $l' \leq l$, where $Km^{l,t}$ is defined from $\phi^{l,t}$. If $\mathcal{A} \subset S^t, t < \infty$ or $\mathcal{A} \subset \{(\mathbf{x}, \lambda^\infty) | \mathbf{x} \in \cup_k S^k\}, t = \infty$ is a decidable set that satisfies (3) and (4) then Theorem 2.2 and Corollary 2.1 hold for $Km^{l,t}$ for $1 \leq l, t \leq \infty$. In order to simplify the argument, in the following discussion, we use Km .

3 Section and relativized randomness

Let P be a computable probability on $X \times Y = \Omega^2$. Let P_X and P_Y be its marginal distributions on X and Y , respectively, i.e., $P_X(x) = P(x, \lambda)$ and $P_Y(y) = P(\lambda, y)$ for $x, y \in S$. Let

$$P(x|y) := \begin{cases} \frac{P(x,y)}{P_Y(y)}, & \text{if } P_Y(y) > 0 \\ 0, & \text{if } P_Y(y) = 0 \end{cases},$$

and

$$P(x|y^\infty) := \lim_{y \rightarrow y^\infty} P(x|y),$$

for $y^\infty \in \Omega$ if the right-hand side exists. For a subset $A \subset X \times Y$ and $y^\infty \in Y$, set

$$A_{y^\infty} := \{x^\infty | (x^\infty, y^\infty) \in A\}.$$

For example, $\mathcal{R}_{y^\infty}^P = \{x^\infty | (x^\infty, y^\infty) \in \mathcal{R}^P\}$. Similarly, for $B \subset S \times S$, set $B_{y^\infty} := \{x | (x, y) \in B, y \sqsubset y^\infty\}$.

Theorem 3.1 ([19]) *If $y^\infty \in \mathcal{R}^{P_Y}$, then $P(x|y^\infty)$ exists for all $x \in S$, and $P(\cdot|y^\infty)$ is a probability measure on $(\Omega, \mathcal{B})$.*

Theorem 3.2 ([19]) *$P(\mathcal{R}_{y^\infty}^P | y^\infty) = 1$ if $y^\infty \in \mathcal{R}^{P_Y}$. $\mathcal{R}_{y^\infty}^P = \emptyset$ if $y^\infty \notin \mathcal{R}^{P_Y}$.*

Corollary 3.1 ([19]) $\mathcal{R}^{P_X} = \bigcup_{y^\infty \in \mathcal{R}^{P_Y}} \mathcal{R}_{y^\infty}^P$.

If $P(\cdot|y^\infty)$ is computable relative to y^∞ , then let $\mathcal{R}^{P(\cdot|y^\infty), y^\infty}$ be the set of random sequences with respect to $P(\cdot|y^\infty)$ relative to y^∞ . In [19], $\{P(\cdot|y^\infty)\}_{y^\infty}$ is called uniformly computable if there is a partial computable A such that $\forall y^\infty \in \mathcal{R}^{P_Y}, x \in S, k \in \mathbb{N} \exists y \sqsubset y^\infty, |P(x|y^\infty) - A(x, y, k)| < 1/k$, i.e., $P(\cdot|y^\infty)$ is uniformly computable relative to all $y^\infty \in \mathcal{R}^{P_Y}$. In [19], it is shown that $\mathcal{R}^{P(\cdot|y^\infty), y^\infty} \subset \mathcal{R}_{y^\infty}^P$, and under uniform computability, $\mathcal{R}^{P(\cdot|y^\infty), y^\infty} = \mathcal{R}_{y^\infty}^P$ for $y^\infty \in \mathcal{R}^{P_Y}$. In the following we show the equivalence without assuming the uniform computability; we only assume that $P(\cdot|y^\infty)$ is computable relative to a given $y^\infty \in \mathcal{R}^{P_Y}$. In order to show $\mathcal{R}^{P(\cdot|y^\infty), y^\infty} \supset \mathcal{R}_{y^\infty}^P$, first we extend a test $U_n^{y^\infty}$ w.r.t. $P(\cdot|y^\infty)$ to a test w.r.t. a finite measure P' on Ω^2 such that the section of the extended test at y^∞ coincide with $U_n^{y^\infty}$ and the total measure of the extended test w.r.t. P' is sufficiently small. Finally by using Markov inequality, we construct a test w.r.t. P .

Theorem 3.3 *Assume that $y^\infty \in \mathcal{R}^{P_Y}$ and $P(\cdot|y^\infty)$ is computable relative to y^∞ , then $\mathcal{R}^{P(\cdot|y^\infty), y^\infty} = \mathcal{R}_{y^\infty}^P$.*

Proof) Fix $y^\infty \in \mathcal{R}^{P_Y}$. Since $P(\cdot|y^\infty)$ is computable relative to y^∞ , there is a partial computable function $A : S \times S \times \mathbb{N} \rightarrow \{q \in \mathbb{Q} | q \geq 0\}$ such that (a1) $\forall x, k \exists y \sqsubset y^\infty, |P(x|y^\infty) - A(x, y, k)| < \frac{1}{k}$ and (a2) if $A(x, y, k)$ is defined then $A(x, y, k) = A(x, z, k)$ for all $y \sqsubset z$. Similarly, let $U^{y^\infty} \subset \mathbb{N} \times S$ be a Martin-Löf test with respect to $P(\cdot|y^\infty)$ relative to y^∞ , i.e., U^{y^∞} is a r.e. set relative to y^∞ , and $P(\tilde{U}_n^{y^\infty} | y^\infty) < 2^{-n}$ for all n , where $U_n^{y^\infty} := \{x | (n, x) \in U^{y^\infty}\}$. Then there is a partial computable function $B : \mathbb{N} \times \mathbb{N} \times S \rightarrow S$ such that

(b1) $\forall n, U_n^{y^\infty} = \{x | \exists i, y \sqsubset y^\infty, B(i, n, y) = x\}$ and (b2) if $B(i, n, y)$ is defined then $B(i, n, y) = B(i, n, z)$ for all $y \sqsubseteq z$.

Let $U_n := \{(x, y) | \exists i, B(i, n, y) = x\}$. Then $U_{n, y^\infty} = U_n^{y^\infty}$. Let $U'_n \subset S \times S$ be a non-overlapping r.e. set such that $\tilde{U}_n = \tilde{U}'_n$. Then $\tilde{U}'_{n, y^\infty} = \tilde{U}_n^{y^\infty}$. Let

$$V_n := \{(x, z, k) \mid (x, y) \in U'_n, y \sqsubseteq z \in S, k \in \mathbb{N}, \frac{1}{k} < \frac{1}{2}A(x, z, k) \text{ or } (k \geq 2^{n+|x|}, A(x, z, k) < \frac{1}{k})\}, \quad (7)$$

$V_n^{X \times Y} := \{(x, y) \mid (x, y, k) \in V_n\}$. Then we have

$$(x, y) \in V_n^{X \times Y} \Rightarrow \forall y \sqsubseteq z, (x, z) \in V_n^{X \times Y}, \quad (8)$$

$$\forall z^\infty \in \Omega, V_{n, z^\infty}^{X \times Y} \text{ is non-overlapping}, \quad (9)$$

$$V_{n, y^\infty}^{X \times Y} = U'_{n, y^\infty}, \quad (10)$$

where (8) follows from (a2); (9) follows from that U'_n is non-overlapping; (10) follows from that: from (a1) and (a2), (i) if $P(x|y^\infty) > 0$ then $\exists y \sqsubset y^\infty, k \forall y \sqsubseteq z, \frac{1}{k} < \frac{1}{2}A(x, z, k)$ and (ii) if $P(x|y^\infty) = 0$ then $\forall k \exists y \sqsubset y^\infty \forall y \sqsubseteq z$ such that $A(x, z, k) < \frac{1}{k}$.

Note that if $\frac{1}{k} < \frac{1}{2}A(x, y, k)$ and $y \sqsubset y^\infty$ then $|P(x|y^\infty) - A(x, y, k)| < \frac{1}{k} < \frac{1}{2}A(x, y, k)$, i.e.,

$$\frac{1}{2}A(x, y, k) < P(x|y^\infty) < \frac{3}{2}A(x, y, k). \quad (11)$$

From V_n , we can construct a r.e. set $W_n \subset S \times S \times \mathbb{N}$ that satisfies (12), (13), (14), (15), and (16) (Lemma 3.1 below):

$$W_n \subset V_n. \quad (12)$$

$$W_n^{X \times Y} \text{ is non-overlapping, where } W_n^{X \times Y} := \{(x, y) | (x, y, k) \in W_n\}. \quad (13)$$

$$(x, y, k), (x, y, k') \in W_n \Rightarrow k = k'. \quad (14)$$

$$\forall z^\infty \in \Omega, \sum_{(x, y, k) \in W_n, y \sqsubseteq z^\infty} A(x, y, k) < 3 \cdot 2^{-n}. \quad (15)$$

$$\tilde{U}_n^{y^\infty} = \tilde{W}_n^{X \times Y}. \quad (16)$$

Let $P'(x, z) := A(x, z, k)P_Y(z)$ for $(x, y, k) \in W_n, y \sqsubseteq z$ and $P'(x, y) := 0$ for (x, y) such that $\Delta(x, y) \cap \tilde{W}_n^{X \times Y} = \emptyset$. Then by (15), $P'(\tilde{W}_n^{X \times Y}) < 3 \cdot 2^{-n}$.

Finally let

$$U_n^{X \times Y} := \{(x, z) \in S \times S \mid (x, y) \in W_n^{X \times Y}, y \sqsubseteq z, \\ P(x, z) < \frac{3}{2}P'(x, z) \text{ or } P(x, z) < 2^{-n-|x|}P_Y(z)\}.$$

Since $W_n^{X \times Y}$ is r.e. and P is computable, we see that $U_n^{X \times Y}$ is a r.e. set. Since $W_n^{X \times Y}$ is non-overlapping, we have $\sum_{(x,y) \in W_n^{X \times Y}} 2^{-|x|}P_Y(y) \leq 1$ and

$$P(\tilde{U}_n^{X \times Y}) < \frac{3}{2}P'(\tilde{W}_n^{X \times Y}) + \sum_{(x,y) \in W_n^{X \times Y}} 2^{-n-|x|}P_Y(y) < \frac{11}{2} \cdot 2^{-n}.$$

From (12), we have $(x, y, k) \in W_n \Rightarrow \frac{1}{k} < \frac{1}{2}A(x, y, k)$ or $k \geq 2^{n+|x|}$, $A(x, y, k) < \frac{1}{k}$. Since $P(x|y) \rightarrow P(x|y^\infty)$ as $y \rightarrow y^\infty$ for $y^\infty \in \mathcal{R}^{P_Y}$ (Theorem 3.1), we have for $(x, y) \in W_n^{X \times Y}$, $y \sqsubseteq y^\infty$ (i) if $\frac{1}{k} < \frac{1}{2}A(x, y, k)$ then from (11), $\exists y \sqsubseteq z \sqsubseteq y^\infty$, $P(x, z) < \frac{3}{2}P'(x, z)$ and (ii) if $k \geq 2^{n+|x|}$, $A(x, y, k) < \frac{1}{k}$ then $\exists y \sqsubseteq z \sqsubseteq y^\infty$, $P(x, z) < 2^{-n-|x|}P_Y(z)$. Thus $\tilde{W}_n^{X \times Y} \subset \tilde{U}_n^{X \times Y}$. Since $\tilde{U}_n^{X \times Y} \subset \tilde{W}_n^{X \times Y}$, from (16), we have

$$\tilde{U}_n^{y^\infty} = \tilde{U}_{n, y^\infty}^{X \times Y}.$$

Since $U^{X \times Y} := \{(n, x, y) \mid (x, y) \in U_n^{X \times Y}\}$ is r.e. and $\sum_n P(\tilde{U}_n^{X \times Y}) < \infty$, we have $\limsup_n \tilde{U}_n^{X \times Y} \subset (\mathcal{R}^P)^c$ and $\mathcal{R}_{y^\infty}^P \subset \mathcal{R}^{P(\cdot|y^\infty), y^\infty}$. The converse inclusion is shown in [19]. $\blacksquare$

Lemma 3.1 *There is a r.e. set W_n that satisfies (12), (14), (15), and (16).*

Proof) We construct a r.e. set $W_n \subset S \times S \times \mathbb{N}$ by induction. Let $W(0) := \emptyset$. Suppose that $W(t-1) \subset V_n$ is finite, $W^{X \times Y}(t-1) := \{(x, z) \mid (x, z, k) \in W(t-1)\}$ is non-overlapping, and

$$\forall z^\infty \in \Omega, \sum_{(x,y,k) \in W(t-1), y \sqsubseteq z^\infty} A(x, y, k) < 3 \cdot 2^{-n}. \quad (17)$$

Since $W(t-1)$ is finite, there is a finite non-overlapping set W^Y such that $\cup_{y \in W^Y} \Delta(y) = \Omega$ and $\sigma\{\Delta(y) \mid y \in W^Y\} = \sigma\{\Delta(y) \mid (x, y, k) \in W(t-1)\}$. Since V_n is a r.e. set, let $v : \mathbb{N} \rightarrow V_n$ be a computable function such that

$v(\mathbb{N}) = V_n$. Let

$$\begin{aligned} w(t) &:= \{(x, z', k) \in S \times S \times \mathbb{N} \mid v(t) = (x, y, k), \\ &\quad \exists z \in W^Y, z' := y \vee z \text{ exists,} \\ &\quad W(t-1) \cup \{(x, z', k)\} \text{ satisfies (17),} \\ &\quad \tilde{W}^{X \times Y}(t-1) \cap \Delta(x, z') = \emptyset\}, \end{aligned}$$

and $W(t) := W(t-1) \cup w(t)$. Let $w^Y := \{z \mid (x, z, k) \in w(t)\}$. Since W^Y is non-overlapping, w^Y is non-overlapping. Hence (i) if $(x, z', k) \in w(t)$ and $z^\infty \in \Delta(z')$ then $\{(x, y, k) \mid (x, y, k) \in W(t), y \sqsubset z^\infty\} = \{(x, y, k) \mid (x, y, k) \in W(t-1), y \sqsubset z^\infty\} \cup \{(x, z', k)\}$ and (ii) if $z^\infty \notin \tilde{w}^Y$ then $\{(x, y, k) \mid (x, y, k) \in W(t), y \sqsubset z^\infty\} = \{(x, y, k) \mid (x, y, k) \in W(t-1), y \sqsubset z^\infty\}$, see Figure 1. Thus (17) holds for $W(t)$. By induction, $W(t)$ is finite and satisfies (17) for all t . Since $W(t-1)$ is finite, we see that $w(t)$ is decidable. Let $W_n := \cup_t W(t)$ then W_n is a r.e. set. Since $\forall t W(t-1) \subset W(t)$, from (17), we have (15). From (8) we have (12). From the last condition of the definition of $w(t)$, we have (13) and (14). From (9), we have $\sum_{x \in V_{n, y^\infty}^{X \times Y}} 2^{-|x|} \leq 1$. Let $V'_{y^\infty} \subset \{(x, y, k) \mid (x, y, k) \in V_n, y \sqsubset y^\infty\}$ such that (i) $(x, y, k), (x, y', k') \in V'_{y^\infty} \Rightarrow y = y', k = k'$ and (ii) $(x, y, k) \in V_n, y \sqsubset y^\infty \Rightarrow \exists y' \sqsubset y^\infty, k', (x, y', k') \in V'_{y^\infty}$. Then for any V'_{y^∞} that satisfies (i) and (ii), from (7) and (11), we have

$$\sum_{(x, y, k) \in V'_{y^\infty}} A(x, y, k) \leq 2P(\tilde{U}_{y^\infty}^n | y^\infty) + \sum_{x \in V_{n, y^\infty}^{X \times Y}} 2^{-n-|x|} < 3 \cdot 2^{-n}.$$

Thus $(x, y, k) \in V_n, y \sqsubset y^\infty \Rightarrow \exists y' \sqsubset y^\infty, k', (x, y', k') \in W_n$ and hence $V_{n, y^\infty}^{X \times Y} \subset W_{n, y^\infty}^{X \times Y}$. From (10) and (12), we have (16). $\blacksquare$

4 Likelihood ratio test

Let P and Q be computable probabilities on Ω . Let

$$r(x) := \begin{cases} \frac{Q(x)}{P(x)}, & \text{if } P(x) > 0 \\ 0, & \text{if } P(x) = 0 \end{cases},$$

for $x \in S$. We see that r is a computable martingale. By the martingale convergence theorem for algorithmically random sequences [19], we have




Figure 1: This figure illustrates a construction of W_n . For example, suppose that $W(t-1) = \{(x_1, y_1, k_1), (x_2, y_2, k_2)\}$, $\alpha = \Delta(x_1, y_1)$, and $\beta = \Delta(x_2, y_2)$ for some t as shown in the figure. W^Y is illustrated by the partition on the Y -axis. If $v(t) = (x_3, y_3, k_3)$ and $\gamma = \Delta(x_3, y_3)$ (the rectangle below α) then γ is divided into $\gamma_1 = \Delta(x_3, y_1)$ and $\gamma_2 = \Delta(x_3, z')$. If $A(x_1, y_1, k_1) + A(x_3, y_1, k_3) < 3 \cdot 2^{-n}$ then $(x_3, y_1, k_3) \in W(t)$, and if $A(x_3, z', k_3) < 3 \cdot 2^{-n}$ then $(x_3, z', k_3) \in W(t)$.

Corollary 4.1 $\mathcal{R}^P \subset \{x^\infty \mid \lim_{x \rightarrow x^\infty} r(x) < \infty\}$.

The following lemma was appeared in [3].

Lemma 4.1 *Let P and Q be computable probabilities on Ω .*

- a) : $\mathcal{R}^P \cap \mathcal{R}^Q = \mathcal{R}^P \cap \{x^\infty \mid 0 < \lim_{x \rightarrow x^\infty} r(x) < \infty\}$.
- b) : $\mathcal{R}^P \cap (\mathcal{R}^Q)^c = \mathcal{R}^P \cap \{x^\infty \mid \lim_{x \rightarrow x^\infty} r(x) = 0\}$.

Proof) a) If $x^\infty \in \mathcal{R}^P \cap \mathcal{R}^Q$ then $P(x) > 0$ and $Q(x) > 0$ for $x \sqsubset x^\infty$. From Corollary 4.1, we have $0 < \lim_{x \rightarrow x^\infty} r(x) < \infty$. Conversely, if $x^\infty \in \mathcal{R}^P \cap \{x^\infty \mid 0 < \lim_{x \rightarrow x^\infty} r(x) < \infty\}$, by Theorem 2.2,

$\sup_{x \sqsubset x^\infty} -\log P(x) - Km(x) < \infty$ and $\sup_{x \sqsubset x^\infty} |-\log Q(x) + \log P(x)| < \infty$. Thus, $\sup_{x \sqsubset x^\infty} -\log Q(x) - Km(x) < \infty$ and we have $x^\infty \in \mathcal{R}^Q$.

b) From a, we have $\mathcal{R}^P \cap (\mathcal{R}^Q)^c = \mathcal{R}^P \cap (\mathcal{R}^P \cap \mathcal{R}^Q)^c = \mathcal{R}^P \cap (\{\lim r = 0\} \cup \{\lim r = \infty\}) = \mathcal{R}^P \cap \{\lim r = 0\}$, where the last equality follows from Corollary 4.1. ■

Remark 3 Let g be an unbounded increasing total-computable function and $\mathcal{A}_{g,n} := \{(x, y) \mid |x| = n, (x, y) \in \mathcal{A}_g\}$, where $\mathcal{A}_g$ is defined in (6). Let

$\mathcal{F}_n := \sigma\{\Delta(x, y) \mid (x, y) \in \mathcal{A}_{g,n}\}$ and $r_n(x^\infty, y^\infty) := \frac{Q(x, y)}{P(x, y)}, (x, y) \in \mathcal{A}_{g,n}$. Then $\{r_n\}$ is martingale with respect to $\{\mathcal{F}_n\}$. If we replace $\lim_{x \rightarrow x^\infty} r(x)$ with $\lim_{(x, y) \rightarrow (x^\infty, y^\infty), (x, y) \in \mathcal{A}_g(x^\infty, y^\infty)} r(x, y)$ in Corollary 4.1 and Lemma 4.1, they hold for computable probabilities on Ω^2 .

Remark 4 In a similar manner with the proof of Lemma 4.1 a), we have $\mathcal{R}^P \cap \mathcal{R}^Q = \mathcal{R}^P \cap \{x^\infty \mid 0 < \inf_{x \sqsubset x^\infty} r(x)\}$. If we replace $\inf_{x \sqsubset x^\infty}$ with $\inf_{(x, y) \in \mathcal{A}_g(x^\infty, y^\infty)}$ for unbounded increasing total-computable g , it holds for computable probabilities on Ω^2 .

4.1 Absolute continuity and mutual singularity

By Lebesgue decomposition theorem, there exists $N \in \mathcal{B}$ such that $P(N) = 0$ and

$$\forall C \in \mathcal{B}, Q(C) = \int_C r(x^\infty) dP + Q(C \cap N). \quad (18)$$

We write (a) $P \perp Q$ if P and Q are mutually singular, i.e., there exist A and B such that $A \cap B = \emptyset$, $P(A) = 1$, and $Q(B) = 1$, and (b) $P \ll Q$ if P is absolutely continuous with respect to Q , i.e., $\forall C \in \mathcal{B} Q(C) = 0 \Rightarrow P(C) = 0$.

Remark 5 By (18), we have (a) $P \perp Q$ iff $P(\{\lim r = 0\}) = 1$, and (b) $P \ll Q$ iff $P(\{\lim r = 0\}) = 0$; for example, see [14].

The following theorem appeared in pp. 103 of [12] without proof.

Theorem 4.1 (Martin-Löf) *Let P and Q be computable probabilities on Ω . Then, $\mathcal{R}^P \cap \mathcal{R}^Q = \emptyset$ iff $P \perp Q$.*

Proof) Since $P(\mathcal{R}^P) = Q(\mathcal{R}^Q) = 1$, only if part follows. Conversely, assume that $P \perp Q$. Let $N := \{x^\infty \mid 0 < \liminf_{x \sqsubset x^\infty} r(x) \leq \limsup_{x \sqsubset x^\infty} r(x) < \infty\}$. By Remark 5, we have $P(N) = Q(N) = 0$. Since $0 < \liminf_{x \sqsubset x^\infty} r(x) \Leftrightarrow 0 < \inf_{x \sqsubset x^\infty} r(x)$ and $\limsup_{x \sqsubset x^\infty} r(x) < \infty \Leftrightarrow \sup_{x \sqsubset x^\infty} r(x) < \infty$, we have

$$\begin{aligned} N &= \{x^\infty \mid 0 < \inf_{x \sqsubset x^\infty} r(x) \leq \sup_{x \sqsubset x^\infty} r(x) < \infty\} \\ &= \bigcup_{a, b \in \mathbb{Q}, 0 < a < b < \infty} \bigcap_{i=1}^{\infty} \tilde{N}_i^{a, b}, \end{aligned}$$

where $N_i^{a,b} = \{x | a \leq r(y) \leq b, \forall y \sqsubseteq x, |x| = i\}$. Since $P(N) = 0$, we have $\lim_i P(\tilde{N}_i^{a,b}) = 0$. Since $(N_i^{a,b})^c \cap \{x | P(x) > 0\}$ is a r.e. set, we can approximate $P(\tilde{N}_i^{a,b})$ from above, and there is a computable function $\alpha(n)$ such that $P(\tilde{N}_{\alpha(n)}^{a,b}) < 2^{-n}$. Thus, $\tilde{N}_{\alpha(n)}^{a,b}$ is a test of P , and hence, $N \subset (\mathcal{R}^P)^c$. From Lemma 4.1 a, we have $\mathcal{R}^P \cap \mathcal{R}^Q = \emptyset$. $\blacksquare$

From Lemma 4.1 b and Remark 5, we have

Lemma 4.2 $\mathcal{R}^P \subset \mathcal{R}^Q \Rightarrow P \ll Q$ for computable probabilities P and Q on Ω .

There is a counter example for the converse implication of the above lemma, see [3]. The above results are related to Kakutani's theorem on product martingale [8, 25], see [7, 23].

4.2 Countable model class

In the following discussion, let $\{P_n\}_{n \in \mathbb{N}}$ be a family of computable probabilities on Ω ; more precisely, we assume that there is a computable function $A : \mathbb{N} \times S \times \mathbb{N} \rightarrow \mathbb{Q}$ such that $|A(n, x, k) - P_n(x)| < 1/k$ for all $n, k \in \mathbb{N}$ and $x \in S$. Note that we cannot set $\{P_n\}_{n \in \mathbb{N}}$ as the entire family of computable probabilities on Ω since it is not a r.e. set. Let α be a computable positive probability on $\mathbb{N}$, i.e., $\forall n \alpha(n) > 0$ and $\sum_n \alpha(n) = 1$. Then, set $P := \sum_n \alpha(n) P_n$. We see that P is a computable probability. The following lemma is a special case (discrete version) of Corollary 3.1

Lemma 4.3 $\mathcal{R}^P = \cup_n \mathcal{R}^{P_n}$.

Proof) Let $P'_Y(y^\infty) := \alpha(n)$ and $P'(x; y^\infty) := P_n(x)$ if $y^\infty = 0^n 10^\infty$ and 0 otherwise, respectively. Let $P'(x, y) := \int_{\Delta(y)} P'(x; y^\infty) dP'_Y$ for $x, y \in S$, then P' is a computable probability on $X \times Y = \Omega^2$. We see that $P'_X(x) = \sum_n \alpha(n) P_n$, $\mathcal{R}^{P'_Y} = \{0^n 10^\infty | n \in \mathbb{N}\}$, and $\mathcal{R}_{y^\infty}^{P'} = \mathcal{R}^{P_n}$ if $y^\infty = 0^n 10^\infty$. Since $\mathcal{R}^{P'_X} = \cup_{y^\infty \in \mathcal{R}^{P'_Y}} \mathcal{R}_{y^\infty}^{P'}$ (Corollary 3.1), we have the lemma. $\blacksquare$

Let β be a computable probability on $\mathbb{N}$ such that 1) $\beta(n) > 0$ if $n \neq n^*$ and $\beta(n^*) = 0$, and 2) $\sum_n \beta(n) = 1$. Then, set

$$P^- := \sum_n \beta(n) P_n.$$

We see that P^- is a computable probability. By Lemma 4.1 and 4.3, we have

Corollary 4.2

$$\mathcal{R}^{P_{n^*}} \cap_{n \neq n^*} (\mathcal{R}^{P_n})^c = (\cup_n \mathcal{R}^{P_n}) \cap \{x^\infty \mid \lim_{x \rightarrow x^\infty} P^-(x)/P_{n^*}(x) = 0\}.$$

Let

$$\hat{n}(x) := \arg \max_n \alpha(n) P_n(x).$$

In [1, 2], it is shown that $\lim_{x \rightarrow x^\infty} P^-(x)/P_{n^*}(x) = 0 \Rightarrow \lim_{x \rightarrow x^\infty} \hat{n}(x) = n^*$. Thus we have

$$\textbf{Corollary 4.3} \quad \mathcal{R}^{P_{n^*}} \cap_{n \neq n^*} (\mathcal{R}^{P_n})^c \subset \{x^\infty \mid \lim_{x \rightarrow x^\infty} \hat{n}(x) = n^*\}.$$

The above corollary shows that if x^∞ is random with respect to $\mathcal{R}^{P_{n^*}}$ and it is not random with respect to other models then $\hat{n}$ classifies its model. Estimation of models by $\hat{n}$ is called MDL model selection, for more details, see [1, 2]. Note that by Theorem 4.1, if $\{P_n\}$ are mutually singular, then $\mathcal{R}^{P_{n^*}} \cap_{n \neq n^*} (\mathcal{R}^{P_n})^c = \mathcal{R}^{P_{n^*}}$, and by Lemma 4.2, if $P_{n^*} \not\ll P^-$, then $\mathcal{R}^{P_{n^*}} \cap_{n \neq n^*} (\mathcal{R}^{P_n})^c \neq \emptyset$.

5 Decomposition of complexity

It can be shown that

$$\sup_{x, y \in S} |Km(x, y) - Km(x|y) - Km(y)| = \infty. \quad (19)$$

The above equation shows that there is a sequence of strings such that the left-hand side of the above equation is unbounded. However, if we restrict strings to an increasing sequence of prefixes of random sequences x^∞, y^∞ with respect to some computable probability and a convergence rate of conditional probability is effective, then we can show that the left-hand-side of (19) is bounded (see Theorem 5.1 below).

Let P be a computable probability on $X \times Y = \Omega^2$. From Theorem 3.1,

$$\forall x, P(x|y) \rightarrow P(x|y^\infty) \text{ as } y \rightarrow y^\infty \in \mathcal{R}^{P_Y}. \quad (20)$$

Observe that

$$P(x, y) > 0, P(x|y^\infty) > 0 \text{ if } (x, y) \sqsubset (x^\infty, y^\infty) \in \mathcal{R}^P. \quad (21)$$

This follows from that $P(x, y) = 0 \Rightarrow (x, y) \sqsubset (x^\infty, y^\infty) \notin \mathcal{R}^P$. If $P(x|y^\infty) = 0$ then from (20), we have $\forall n \exists y \sqsubset y^\infty, P(x|y) < 2^{-n}$. Since $U_n := \{(x, y) \mid P(x, y) < 2^{-n} P_Y(y)\}$ is a test of P , we have $(x^\infty, y^\infty) \in \cap_n \tilde{U}_n$ if $P(x|y^\infty) = 0$.

If $(x^\infty, y^\infty) \in \mathcal{R}^P$ then from (20) and (21), we have

$$\forall x \sqsubset x^\infty, f > 0 \exists N \forall y \sqsubset y^\infty, N \leq |y| \Rightarrow \left| \frac{P(x|y)}{P(x|y^\infty)} - 1 \right| < f.$$

By letting f be a function of $|x|$, we have for any $f : \mathbb{N} \rightarrow \{q \in \mathbb{Q} | q > 0\}$, there is $g : \mathbb{N} \rightarrow \mathbb{N} \cup \{0\}$ such that

$$\forall (x, y) \sqsubset (x^\infty, y^\infty), g(|x|) \leq |y| \Rightarrow \left| \frac{P(x|y)}{P(x|y^\infty)} - 1 \right| < f(|x|). \quad (22)$$

In the above, g depends on f and (x^∞, y^∞) . We say that the conditional probability $P(\cdot|y^\infty)$ is $f, (x^\infty, y^\infty)$ *effectively converges* if there is a total-computable monotonically increasing g in (22), where we allow that g is bounded, see Remark 6. g is called effective convergence rate function.

Lemma 5.1 *Let P be a computable probability on $X \times Y = \Omega^2$ and $(x^\infty, y^\infty) \in \mathcal{R}^P$. Let $f : \mathbb{N} \rightarrow \{q \in \mathbb{Q} | 0 < q < 1\}$ such that $\sum_n f(n) < \infty$. Assume that $P(\cdot|y^\infty)$ is $f, (x^\infty, y^\infty)$ effectively converges. Let g be an effective convergence rate function. Then there is a computable monotone function $e : (S \cup \Omega)^2 \rightarrow S \cup \Omega$ such that*

$$\begin{aligned} \exists c \exists p^\infty \in \Omega \forall (x, y) \sqsubset (x^\infty, y^\infty) \exists p \sqsubset p^\infty, \\ g(|x|) = |y| \Rightarrow x \sqsubseteq e(p, y), |p| \leq -\log P(x|y) + c. \end{aligned} \quad (23)$$

$$\exists c \forall (x, y) \sqsubset (x^\infty, y^\infty), g(|x|) = |y| \Rightarrow Km(x|y) \leq -\log P(x|y) + c. \quad (24)$$

Proof) Let

$$\begin{aligned} P'(0|y^\infty) &:= P(0|y) \text{ for } |y| = g(1), y \sqsubset y^\infty, \\ P'(1|y^\infty) &:= 1 - P'(0|y^\infty), \end{aligned} \quad (25)$$

and for $x \in S$

$$\begin{aligned} P'(x0|y^\infty) &:= P'(x|y^\infty) \frac{P(x0|y)}{P(x|y)} \text{ if } P(x|y) > 0 \text{ for } |y| = g(|x| + 1), y \sqsubset y^\infty, \\ P'(x1|y^\infty) &:= P'(x|y^\infty) - P'(x0|y^\infty). \end{aligned} \quad (26)$$

Since $P(x|y) > 0 \Leftrightarrow \forall (x', y') \sqsubseteq (x, y), P(x'|y') > 0$ and g is computable, we see that there is a partial computable $A : S \times S \times \mathbb{N} \rightarrow \mathbb{Q}$ such that

$$\forall y^\infty \forall x, y, k, |P'(x|y^\infty) - A(x, y, k)| \leq \frac{1}{k} \text{ if } y \sqsubset y^\infty, g(|x|) = |y|, P(x|y) > 0. \quad (27)$$

Let $D := \{(x, y) | g(|x|) = |y|, P(x|y) > 0\}$. From (27), we can construct a family of half-open intervals $V_{(x,y)} \subset [0, 1]$, $(x, y) \in D$ such that 1) the endpoints of $V_{(x,y)}$ are computable with arbitrary precision from $(x, y) \in D$ and $|V_{(x,y)}| = P'(x|y^\infty)$, and 2) if $(x, y), (x', y') \in D$, and y and y' are comparable, then i) $x \sqsubseteq x' \Rightarrow V_{(x',y')} \subset V_{(x,y)}$, and ii) $\Delta(x) \cap \Delta(x') = \emptyset \Rightarrow V_{(x,y)} \cap V_{(x',y')} = \emptyset$. Let $F := \{(s, y, x) | I_s \subset V_{(x,y)}, (x, y) \in D\} \cup \{(s, y, \lambda) | s, y \in S\}$. Then F is r.e. and satisfies (1). Let e be the monotone function defined by F . Then

$$\forall y^\infty \exists c \forall x, y, Km_e(x|y) \leq -\log P'(x|y^\infty) + c \text{ if } y \sqsubset y^\infty, g(|x|) = |y|, P(x|y) > 0. \quad (28)$$

By replacing $P(x|y)$ in (25) with $P(x|y^\infty)$, from (22), we have for $|x| = 1$,

$$(1 - f(1)) \leq \frac{P'(x|y^\infty)}{P(x|y^\infty)} \leq (1 + f(1)).$$

Similarly, by replacing $P(xz|y)$ and $P(x|y)$ in (26) with $P(xz|y^\infty)$ and $P(x|y^\infty)$ respectively, from (22), we have for $1 \leq |x|, |z| = 1$,

$$\frac{1 - f(|x| + 1)}{1 + f(|x|)} \frac{P'(x|y^\infty)}{P(x|y^\infty)} \leq \frac{P'(xz|y^\infty)}{P(xz|y^\infty)} \leq \frac{P'(x|y^\infty)}{P(x|y^\infty)} \frac{1 + f(|x| + 1)}{1 - f(|x|)}.$$

Therefore we have

$$\frac{\prod_{n=1}^{|x|} (1 - f(n))}{\prod_{n=1}^{|x|-1} (1 + f(n))} \leq \frac{P'(x|y^\infty)}{P(x|y^\infty)} \leq \frac{\prod_{n=1}^{|x|} (1 + f(n))}{\prod_{n=1}^{|x|-1} (1 - f(n))} \text{ if } P(x|y^\infty) > 0.$$

Since $0 < \prod_{n=1}^\infty (1 - f(n)) \leq \prod_{n=1}^\infty (1 + f(n)) < \infty$ if $\sum_n f(n) < \infty$ and $0 < f < 1$, from (28), we have the lemma. $\blacksquare$

Theorem 5.1 *Let P be a computable probability on $X \times Y = \Omega^2$ and $(x^\infty, y^\infty) \in \mathcal{R}^P$. Let $f : \mathbb{N} \rightarrow \{q \in \mathbb{Q} | 0 < q < 1\}$ such that $\sum_n f(n) < \infty$. Assume that $P(\cdot|y^\infty)$ is $f, (x^\infty, y^\infty)$ effectively converges. Let g be an effective convergence rate function. Then*

$$\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |Km(x|y) + \log P(x|y)| < \infty, \quad (29)$$

$$\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |Km(x, y) - Km(x|y) - Km(y)| < \infty, \quad (30)$$

where $\mathcal{A}_g$ is defined in (6). In addition, if $P(\cdot|y^\infty)$ is computable relative to y^∞ , then

$$\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} Km(x|y) - Km(x|y^\infty) < \infty. \quad (31)$$

Proof) From Corollary 2.1, if $(x^\infty, y^\infty) \in \mathcal{R}^P$ then

$$\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |\log P(x, y) + Km(x, y)| < \infty, \sup_{y \sqsubset y^\infty} |\log P_Y(y) + Km(y)| < \infty, \quad (32)$$

$$\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} -\log P(x|y) - Km(x|y) < \infty. \quad (33)$$

From (24) and (33), we have (29). From (32) and (29), we have (30).

If $P(\cdot|y^\infty)$ is computable relative to y^∞ , Theorem 3.3 holds, i.e., $(x^\infty, y^\infty) \in \mathcal{R}^P$ iff $x^\infty \in \mathcal{R}^{P(\cdot|y^\infty), y^\infty}$, $y^\infty \in \mathcal{R}^{P_Y}$. By relativized version of Levin-Schnorr theorem, we have

$$\sup_{x \sqsubset x^\infty} -\log P(x|y^\infty) - Km(x|y^\infty) < \infty,$$

for $(x^\infty, y^\infty) \in \mathcal{R}^P$. Since $Km(x|y^\infty) \leq Km(x|y)$ for $y \sqsubset y^\infty$, from (22) and (29), we have (31). $\blacksquare$

Example 2 Let P' be a computable probability on Ω . For $x = x_1 \cdots x_n, y = y_1 \cdots y_m \in S$, let

$$\Delta(x \oplus y) := \{z_1 z_2 \cdots \in \Omega \mid \begin{array}{l} z_i = x_i \text{ if } i \text{ is odd and } i \leq n, \\ z_i = y_i \text{ if } i \text{ is even and } i \leq m \end{array}\},$$

$$P(x, y) := P'(\Delta(x \oplus y)).$$

Then P is a computable probability on $X \times Y = \Omega^2$, i.e., X and Y are the spaces of odd and even coordinates, respectively. For $x^\infty = x_1 x_2 \cdots, y^\infty = y_1 y_2 \cdots$, let

$$x^\infty \oplus y^\infty := x_1 y_1 x_2 y_2 \cdots \in \Omega,$$

then

$$(x^\infty, y^\infty) \in \mathcal{R}^P \iff x^\infty \oplus y^\infty \in \mathcal{R}^{P'}.$$

From Theorem 3.3, if the conditional probability is computable relative to $y^\infty \in \mathcal{R}^{P_Y}$ then $x^\infty \oplus y^\infty$ is random with respect to P' iff y^∞ is random and x^∞ is random with respect to the conditional probability at y^∞ . Let P' be a computable first order Markov process, i.e., $P'(z_1 \cdots z_n) = p_{z_1} \prod_{i=2}^n p_{z_{i-1}, z_i}$, where $\forall i, j \in \{0, 1\}, 0 \leq p_i, p_{i,j} \leq 1, \sum_i p_i = 1, \sum_j p_{i,j} = 1$. We see that $P(x|y^\infty) = P(x|y_1 \cdots y_{|x|})$. Thus $g(n) = n$ satisfies (22) for any f and Theorem 5.1 holds.

Remark 6 In Lemma 5.1 and Theorem 5.1, g need not be unbounded if (22) hold. For example if $P := P_X P_Y$ then $g := 0$ satisfies (22) for any f .

5.1 Independence

We show some equivalent conditions for independence of two individual sequences. The following result shows that if (x^∞, y^∞) is random with respect to some computable probability (in [13] such a sequence is called *natural*), then we can represent independence of (x^∞, y^∞) in terms of complexity.

Corollary 5.1 *Let P be a computable probability on Ω^2 and $(x^\infty, y^\infty) \in \mathcal{R}^P$. Assume that $P(\cdot|y^\infty)$ is computable relative to y^∞ and $f, (x^\infty, y^\infty)$ effectively converges for $f = 1$. Let Q be a computable probability such that $\forall x, y, Q(x, y) := P_X(x)P_Y(y)$. The following statements are equivalent:*

- a) $(x^\infty, y^\infty) \in \mathcal{R}^Q$.
- b) for any unbounded computable increasing g ,
 $\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |Km(x, y) - Km(x) - Km(y)| < \infty$.
- c) $\sup_{x \sqsubseteq x^\infty} Km(x) - Km(x|y^\infty) < \infty$.

Proof) a $\Rightarrow$ b: Every increasing computable g satisfies (22) for Q . From Theorem 5.1, if $(x^\infty, y^\infty) \in \mathcal{R}^Q$ then $\sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |Km(x|y) + \log P_X(x)| < \infty$, $\sup_{x \sqsubseteq x^\infty} |Km(x) + \log P_X(x)| < \infty$, and (30) holds. Thus we have b.

b $\Rightarrow$ a: Let g be unbounded computable increasing function. Since $\mathcal{R}^P \subset \mathcal{R}^{P_X} \times \mathcal{R}^{P_Y}$,

$$\begin{aligned} (x^\infty, y^\infty) \in \mathcal{R}^P \Rightarrow & \sup_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} |Km(x, y) + \log P(x, y)| < \infty, \\ & \sup_{x \sqsubseteq x^\infty} |Km(x) + \log P_X(x)| < \infty, \\ & \sup_{y \sqsubseteq y^\infty} |Km(y) + \log P_Y(y)| < \infty. \end{aligned}$$

We have $0 < \inf_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} \frac{Q(x,y)}{P(x,y)}$. From Lemma 4.1 (see Remark 4), we have a.

a $\Rightarrow$ c: Since $g := 0$ satisfies (22) for Q , from Theorem 5.1, we have c, see Remark 6.

c $\Rightarrow$ a: Let g be an unbounded effective convergence rate function for $P(\cdot|y^\infty)$, $f = 1$, and $(x^\infty, y^\infty) \in \mathcal{R}^P$. Then we have $\frac{P_X(x)}{2P(x|y^\infty)} \leq \frac{P_X(x)}{P(x|y)} = \frac{Q(x,y)}{P(x,y)}$ for $(x, y) \in \mathcal{A}_g(x^\infty, y^\infty)$. From Theorem 3.3 and Levin-Schnorr theorem, we have

$\sup_{x \sqsubset x^\infty} |Km(x|y^\infty) + \log P(x|y^\infty)| < \infty$ and $\sup_{x \sqsubset x^\infty} |Km(x) + \log P_X(x)| < \infty$. From the statement c), we have $0 < \inf_{(x,y) \in \mathcal{A}_g(x^\infty, y^\infty)} \frac{Q(x,y)}{P(x,y)}$. From Lemma 4.1 (see Remark 4), we have a. $\blacksquare$

Note that $\mathcal{R}^P \cap \mathcal{R}^Q \neq \emptyset$ iff P and Q are not mutually singular (Theorem 4.1) iff $P(\lim r > 0) > 0$ (Remark 5).

6 Bayesian statistics

Let P be a computable probability on $X \times Y$ and P_X, P_Y be its marginal distributions as before. In Bayesian statistical terminology, if X is a sample space, then P_X is called mixture distribution, and if Y is a parameter space, then P_Y is called prior distribution. We show that section of random set satisfies many theorem of Bayesian statistics, see also [19], and it is natural as a definition of random set with respect to conditional probability from Bayesian statistical point of view.

6.1 Consistency of posterior distribution

We show a consistency of posterior distribution for algorithmically random sequences. We see that the classification of random sets by likelihood ratio test (see Section 4) plays an important role in this section.

Theorem 6.1 *Let P be a computable probability on $X \times Y = \Omega^2$. The following six statements are equivalent:*

- a) $P(\cdot|y) \perp P(\cdot|z)$ if $\Delta(y) \cap \Delta(z) = \emptyset, P_Y(y) > 0, P_Y(z) > 0$ for $y, z \in S$.
- b) $\mathcal{R}^{P(\cdot|y)} \cap \mathcal{R}^{P(\cdot|z)} = \emptyset$ if $\Delta(y) \cap \Delta(z) = \emptyset, P_Y(y) > 0, P_Y(z) > 0$ for $y, z \in S$.
- c) $P_{Y|X}(\cdot|x)$ converges weakly to I_{y^∞} as $x \rightarrow x^\infty$ for $(x^\infty, y^\infty) \in \mathcal{R}^P$, where I_{y^∞} is the distribution that has probability of 1 at y^∞ .
- d) $\mathcal{R}_{y^\infty}^P \cap \mathcal{R}_{z^\infty}^P = \emptyset$ if $y^\infty \neq z^\infty$.
- e) There exists a surjective function $f : \mathcal{R}^{P_X} \rightarrow \mathcal{R}^{P_Y}$ such that $f(x^\infty) = y^\infty$ for $(x^\infty, y^\infty) \in \mathcal{R}^P$.
- f) There exists $f : X \rightarrow Y$ and $Y' \subset Y$ such that $P_Y(Y') = 1$ and $f = y^\infty, P(\cdot|y^\infty) - a.s.$ for $y^\infty \in Y'$.

Proof) a $\Leftrightarrow$ b follows from Theorem 4.1.

b $\Rightarrow$ c : If $(x^\infty, y^\infty) \in \mathcal{R}^P$, then $x^\infty \in \mathcal{R}^{P(\cdot|y)}$ and $P_Y(y) > 0$ for $y \sqsubset y^\infty$. If

$\Delta(y) \cap \Delta(z) = \emptyset$ and $P_Y(z) > 0$, then from the statement b, $x^\infty \notin \mathcal{R}^{P(\cdot|z)}$. If $P_Y(z) > 0$ then from Lemma 4.1, we have $\lim_{x \rightarrow x^\infty} P(x|z)/P(x|y) = 0$, and

$$\lim_{x \rightarrow x^\infty} \frac{P(x|z)}{P(x|y)} = 0 \Leftrightarrow \lim_{x \rightarrow x^\infty} \frac{P(x, z)}{P(x, y)} = 0 \Leftrightarrow \lim_{x \rightarrow x^\infty} \frac{P_{Y|X}(z|x)}{P_{Y|X}(y|x)} = 0. \quad (34)$$

If $P_Y(z) = 0$ then the last equation in (34) holds. Hence the last equation in (34) holds for all z and we see that the posterior distribution $P_{Y|X}(\cdot|x)$ converges weakly to I_{y^∞} .

c $\Rightarrow$ d : obvious.

d $\Rightarrow$ e : Since $\mathcal{R}_{y^\infty}^P \cap \mathcal{R}_{z^\infty}^P = \emptyset$ for $y^\infty \neq z^\infty$, we can define a function $f : X \rightarrow Y$ such that $f(x^\infty) = y^\infty$ for $x^\infty \in \mathcal{R}_{y^\infty}^P$. From Corollary 3.1, we have e, see Figure 2.

e $\Rightarrow$ f : By Theorem 3.2, we have f.

f $\Rightarrow$ a : Let $A_{y^\infty} := \{x^\infty | f(x^\infty) = y^\infty\}$. Then, $A_{y^\infty} \cap A_{z^\infty} = \emptyset$ for $y^\infty \neq z^\infty$ and $P(A_{y^\infty} | y^\infty) = 1$ for $y^\infty \in Y'$. Thus, $(\cup_{y^\infty \in \Delta(y)} A_{y^\infty}) \cap (\cup_{y^\infty \in \Delta(z)} A_{y^\infty}) = \emptyset$ for $\Delta(y) \cap \Delta(z) = \emptyset$ and $P(\cup_{y^\infty \in \Delta(y)} A_{y^\infty} | y) = P(\cup_{y^\infty \in \Delta(z)} A_{y^\infty} | z) = 1$, which shows a. $\blacksquare$

Usually, consistency of posterior distribution is derived from f, see [6]. Note that the statements a and f do not contain algorithmic notion.

Example 3 Let $\{P(\cdot; y^\infty)\}_{y^\infty \in Y}$ be the parametric model of Bernoulli process, i.e., $P(x; y^\infty) := r(y^\infty)^{\sum_{i=1}^n x_i} (1 - r(y^\infty))^{n - \sum_{i=1}^n x_i}$ where $x = x_1 \cdots x_n$, $y^\infty = y_1 y_2 \cdots$, and $r(y^\infty) := \sum_i y_i 2^{-i}$. Let P_Y be a computable probability on Ω and $P(x, y) := \int_{\Delta(y)} P(x; y^\infty) dP_Y$ for $x, y \in S$. Then P is a computable probability on Ω^2 . By the law of large numbers, f (and all the statements) are satisfied. Note that the conditional probability $P(\cdot | y^\infty)$ is defined by P , see Section 4 in [19]. In general, it is possible that $P(\cdot | y^\infty) \neq P(\cdot; y^\infty)$ at y^∞ of a null set.

6.2 Algorithmically best estimator

We study asymptotic theory of estimation for individual samples and parameters from algorithmic point of view.

Suppose that one of the statement of Theorem 6.1 holds. Then from the statement c, we have $P(y|x^\infty) = 1$ for $y \sqsubset y^\infty, (x^\infty, y^\infty) \in \mathcal{R}^P$. Since $P(y|x) \rightarrow P(y|x^\infty)$ as $x \rightarrow x^\infty$ if $x^\infty \in \mathcal{R}^{P_X}$, we have $\forall \epsilon > 0, y \sqsubset y^\infty, \exists x \sqsubset$




Figure 2: $f : \mathcal{R}^{P_X} \rightarrow \mathcal{R}^{P_Y}$

$x^\infty, P(y|x) > 1 - \epsilon$. In particular there is an increasing h such that $\forall \epsilon, y \sqsubseteq y^\infty, x \sqsubseteq x^\infty, |x| \geq h(|y|) \Rightarrow P(y|x) > 1 - \epsilon$. Roughly speaking, the following theorem shows that if this happen then y is estimated from x of size h and if $P(y|x)$ goes to 0 then we cannot estimate y from sample size h .

Theorem 6.2 *Let P be a computable probability on $X \times Y = \Omega^2$. Let $h : \mathbb{N} \rightarrow \mathbb{N}$ be an increasing computable function and $\mathcal{A} := \{(x, y) | |x| = h(|y|)\}$. For each (x^∞, y^∞) we have:*

- a) *If $\inf_{(x,y) \in \mathcal{A}(x^\infty, y^\infty)} P(y|x) > 0$, then there is a computable function ρ such that $y = \rho(x)$ for infinitely many $(x, y) \in \mathcal{A}(x^\infty, y^\infty)$, where ρ need not be monotone.*
- b) *Let $f : \mathbb{N} \rightarrow \{q \in \mathbb{Q} | 0 < q < 1\}$ such that $\sum_n f(n) < \infty$. Assume that $P(\cdot | x^\infty)$ effectively converges for f and $(x^\infty, y^\infty) \in \mathcal{R}^P$, i.e., there is a total computable increasing $h : \mathbb{N} \rightarrow \mathbb{N}$ such that*

$$|x| = h(|y|) \Rightarrow \left| \frac{P(y|x)}{P(y|x^\infty)} - 1 \right| < f(|y|).$$

If $\inf_{(x,y) \in \mathcal{A}(x^\infty, y^\infty)} P(y|x) > 0$ then there is a computable monotone function ρ such that $\forall (x, y) \in \mathcal{A}(x^\infty, y^\infty), y \sqsubseteq \rho(x)$.

- c) *If $(x^\infty, y^\infty) \in \mathcal{R}^P$ and $\inf_{(x,y) \in \mathcal{A}(x^\infty, y^\infty)} P(y|x) = 0$, then there is no computable monotone function ρ such that $\forall (x, y) \in \mathcal{A}(x^\infty, y^\infty), y \sqsubseteq \rho(x)$.*

Proof) a) By applying Shannon-Fano-Elias coding to $P(\cdot | x)$ on the finite partition $\{y | |y| = h^{-1}(|x|)\}$, we can construct a computable function e and

a program $p \in S$ such that $e(p, x) = y$ and $|p| = \lceil -\log P(y|x) \rceil + 1$. Here, e need not be a monotone function. Since $|p| < \infty$ as $x \rightarrow x^\infty$, there is a p_0 such that $e(p_0, x) = y$ for infinitely many prefix x of x^∞ . Thus, $\rho(x) := e(p_0, x)$ satisfies a.

b) From (23), there is a computable monotone function e and $p \in S$ such that $\forall(x, y) \sqsubset \mathcal{A}(x^\infty, y^\infty), y \sqsubseteq e(p, x)$. Let $\rho(x) := e(p, x)$ then ρ satisfies b.

c) As in the same way of (33), we have $\sup_{(x, y) \in \mathcal{A}(x^\infty, y^\infty)} -\log P(y|x) - Km(y|x) < \infty$. Since $\sup_{(x, y) \in \mathcal{A}(x^\infty, y^\infty)} -\log P(y|x) = \infty$, we have $\sup_{(x, y) \in \mathcal{A}(x^\infty, y^\infty)} Km(y|x) = \infty$. If there is a computable monotone function ρ such that $\forall(x, y) \in \mathcal{A}(x^\infty, y^\infty), y \sqsubseteq \rho(x)$ then $\sup_{(x, y) \in \mathcal{A}(x^\infty, y^\infty)} Km(y|x) < \infty$, which is a contradiction. $\blacksquare$

By definition, we have

$$-\log P(y|x) = -\log \int_{\Delta(y)} P(x|y^\infty) dP_Y(y^\infty) + \log \int_Y P(x|y^\infty) dP_Y(y^\infty). \quad (35)$$

Let P_Y be a Lebesgue absolutely continuous measure. Let $\hat{y}$ be the maximum likelihood estimator. By using Laplace approximation with suitable conditions, if $\hat{y} \in \Delta(y)$ and $h^{-1}(|x|) \approx \frac{1}{2} \log |x|$, then the right-hand-side of (35) is asymptotically bounded, for example see [1], and we have $\inf_{x \sqsubset x^\infty} P(y|x) > 0$, where $|y| = h^{-1}(|x|)$. Thus, by Theorem 6.2 a, we can compute initial $\lceil \frac{1}{2} \log |x| \rceil$ -bits of y^∞ from x infinitely many times, which is an algorithmic version of a well known result in statistics: $|y^\infty - \hat{y}| = O(1/\sqrt{n})$.

Let $h^{-1}(\cdot)$ be a large order function such that $\inf_{x \sqsubset x^\infty} P(y|x) = 0$ for $|y| = h^{-1}(|x|)$; for example, set $h^{-1}(|x|) = \lceil \log |x| \rceil$. By Theorem 6.2 c, there is no monotone computable function that computes initial $h^{-1}(|x|)$ -bits of y^∞ for all $x \sqsubset x^\infty$. If such a function exists, then y^∞ is not random with respect to P_Y and the Lebesgue measure of such parameters is 0. On the other hand, it is known that the set of parameters that are estimated within $o(1/\sqrt{n})$ accuracy has Lebesgue measure 0 [4].

Theorem 6.2 shows a relation between the redundancy of universal coding and parameter estimation; as in [18], if we set P_Y to be a singular prior, we have $\inf_{x \sqsubset x^\infty} P(y|x) > 0$ for a large order h^{-1} . In such a case we have a super-efficient estimator.

Acknowledgement

The author thanks Prof. Teturo Kamae (Matsuyama Univ.), Prof. Akio Fujiwara, and Masahiro Nakamura (Osaka Univ.) for discussions and comments

to earlier versions of the paper. The author also thanks anonymous referees for valuable comments, which helped to improve the paper.

References

- [1] A. Barron, J. Rissanen, and B. Yu. The minimum description length principle in coding and modeling. *IEEE Trans. Inform. Theory*, 44(6):2743–2760, 1998.
- [2] A. R. Barron. *Logically smooth density estimation*. PhD thesis, Stanford Univ., 1985.
- [3] L. Bienvenu and W. Merkle. Effective randomness for computable probability measures. *Electron. Notes Theor. Comput. Sci.*, 167:117–130, 2007.
- [4] L. Le Cam. On some asymptotic properties of maximum likelihood estimates and related Bayes estimates. *University of California Publications in Statistics*, 1:277–330, 1953.
- [5] A. Chernov and M. Hutter. Monotone conditional complexity bounds on future prediction errors. In *ALT 2005*, volume 3734 of *LNAI*, pages 414–428. Springer, 2005.
- [6] J. L. Doob. Application of the theory of martingales. In *Le Calcul des Probabilités et ses Applications. Colloq. Intern. du CNRS*, pages 22–28, Paris, 1948.
- [7] A. Fujiwara. Randomness criteria in terms of α -divergences. *IEEE Trans. Inform. Theory*, 54(3):1252–1261, 2008.
- [8] S. Kakutani. On equivalence of infinite product measures. *Ann. Math.*, 49:214–224, 1948.
- [9] L. A. Levin. On the notion of a random sequence. *Soviet. Math. Dokl.*, 14(5):1413–1416, 1973.
- [10] M. Li and P. Vitányi. *An introduction to Kolmogorov complexity and Its applications*. Springer, New York, second edition, 1997.
- [11] P. Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–609, 1966.
- [12] P. Martin-Löf. *Notes on constructive mathematics*. Almqvist & Wiksell, Stockholm, 1968.

- [13] A. A. Muchnik, A. L. Semenov, and V. A. Uspensky. Mathematical metaphysics of randomness. *Theoret. Comput. Sci.*, 207:263–317, 1998.
- [14] J. Neveu. *Discrete-Parameter Martingales*. North-Holland, Amsterdam, 1975.
- [15] C. P. Schnorr. Process complexity and effective random tests. *J. Comp. Sys. Sci.*, 7:376–388, 1973.
- [16] C. P. Schnorr. A survey of the theory of random sequences. In Butts and Hintikka, editors, *Basic problems in Methodology and Linguistics*, pages 193–211. Reidel, Dordrecht, 1977.
- [17] A. Kh. Shen. On relations between different algorithmic definitions of randomness. *Soviet Math. Dokl.*, 38(2):316–319, 1989.
- [18] H. Takahashi. Redundancy of universal coding, Kolmogorov complexity, and Hausdorff dimension. *IEEE Trans. Inform. Theory*, 50(11):2727–2736, 2004.
- [19] H. Takahashi. On a definition of random sequences with respect to conditional probability. *Inform. and Compt.*, 206:1375–1382, 2008.
- [20] V. A. Uspenskii, A. L. Semenov, and A. Kh. Shen. Can an individual sequence of zeros and ones be random? *Russian Math. Surveys*, 45(1):121–189, 1990.
- [21] V. A. Uspensky and A. Shen. Relation between varieties of Kolmogorov complexities. *Theory Comput. Syst.*, 29:271–292, 1996.
- [22] M. van Lambalgen. *Random sequences*. PhD thesis, Universiteit van Amsterdam, 1987.
- [23] V. G. Vovk. On a randomness criterion. *Soviet Math. Dokl.*, 35(3):656–660, 1987.
- [24] V. V. V'yugin. Ergodic theorems for individual random sequences. *Theor. Comp. Sci.*, 207:343–361, 1998.
- [25] D. Williams. *Probability with Martingale*. Cambridge university press, Cambridge, 1991.